

Neosporin® Polymyxin B-Bacitracin-Neomycin

REFERENCES

1. Zimmerman B: Intravenous tubing for parenteral therapy. *Science* 101:567-568, 1945
2. Meyers L: Intravenous catheterization. *Amer J Nurs* 45:930-931, 1945
3. Duffy BJ Jr: Clinical use of polyethylene tubing for intravenous therapy—Report on 72 cases. *Ann Surg* 130:929-936, 1949
4. Bonner CD: Experience with plastic tubing in prolonged intravenous therapy. *New Eng J Med* 245:97-98, 1951
5. Bansmer G, Keith D, Tesluk H: Complications following use of indwelling catheters of inferior vena cava. *JAMA* 167:1606-1611, 1958
6. Cheney FW Jr, Lincoln JR: Phlebitis from plastic intravenous catheters. *Anesthesiology* 25:650-652, 1964
7. Moncrief JA: Femoral catheters. *Ann Surg* 147:166-172, 1958
8. O'Neill JA Jr, Pruitt BA Jr, Foley FD, et al: Suppurative thrombophlebitis—A lethal complication of intravenous therapy. *J Trauma* 8:256-267, 1968
9. Bogen JE: Local complications of 167 patients with indwelling venous catheters. *Surg Gynec Obstet* 110:112-114, 1960
10. Druskin MS, Siegel PD: Bacterial contamination of indwelling intravenous polyethylene catheters. *JAMA* 185:966-968, 1963
11. Phillips RW, Eyre JD Jr: Septic thrombophlebitis with septicemia—Report of three cases due to *Staphylococcus aureus* infection after the intravenous use of polyethylene catheters for parenteral therapy. *New Eng J Med* 259:729-731, 1958
12. Smits H, Freedman LR: Prolonged venous catheterization as a cause of sepsis. *New Eng J Med* 276:1229-1233, 1967
13. Wellman KF, Reinhard A, Salazar EP: Polyethylene catheter embolism—Review of the literature and report of a case with associated fatal tricuspid and systemic candidiasis. *Circulation* 37:380-392, 1968
14. Deering RB, Stemmer EA, Connolly JE: Complications of indwelling venous catheters, with particular reference to catheter embolism. *Amer J Surg* 114:259-266, 1967
15. Brown CA, Kent A: Perforation of right ventricle by polyethylene catheter. *Southern Med J* 49:466-467, 1956
16. Turner DD, Sommers SC: Accidental passage of a polyethylene catheter from cubital vein to right atrium—Report of a fatal case. *New Eng J Med* 251:744-745, 1954
17. Breed RS, Murray EGD, Smith NR: *Bergey's Manual of Determinative Bacteriology* (7th ed.). Baltimore, Williams and Wilkins Co, 1957
18. Isenberg HD, Berkman JJ: Recent practices in diagnostic bacteriology. *In Progress in Clinical Pathology*. New York, Grune and Stratton, Inc, 1966
19. Bailey WR, Scott EG: *Diagnostic Microbiology—A Textbook for the Isolation and Identification of Pathogenic Microorganisms*. St. Louis, Missouri, V Mosby Co, 1966
20. Bentley DW, Lepper MH: Septicemia related to indwelling venous catheter. *JAMA* 206:1749-1752, 1968
21. Moran JM, Atwood RP, Rowe MI: A clinical and bacteriologic study of infections associated with venous cutdowns. *New Eng J Med* 272:554-560, 1965
22. Collins RN, Braun PA, Zinner SH, et al: Risk of local and systemic infection with polyethylene intravenous catheters. *New Eng J Med* 279:340-343, 1968
23. Cluff LE, Reynolds RC, Page DL, et al: Staphylococcal bacteremia and altered host resistance. *Ann Intern Med* 69:859-873, 1968
24. Hassall JE, Rountree PM: Staphylococcal septicemia. *Lancet* 1:213-217, 1959
25. Andriole VT, Kravetz HM, Roberts WC, et al: Candida endocarditis—Clinical and pathologic studies. *Amer J Med* 32:251-285, 1962
26. Louria DB, Stiff DP, Bennett B: Disseminated moniliasis in the adult. *Medicine* 41:307-337, 1962
27. Michaels L, Ruebner B: Growth of bacteria in intravenous infusion fluids. *Lancet* 1:772-774, 1953
28. Michaels L: Cannula sepsis. *New Eng J Med* 277:433, 1967
29. Worms R: Les complications septicemiques des catheterisations intraveineuses prolongees. *Bull Soc Hop Paris* 117:103-106, Jan 1966
30. Morse LJ, Williams HL, Grenn FP Jr, et al: Septicemia due to *Klebsiella pneumoniae* originating from a hand-cream dispenser. *New Eng J Med* 277:472-473, 1967
31. Norden CW: Application of antibiotic ointment to the site of venous catheterization—A controlled trial. *J Infect Dis* 120:611-615, 1969
32. Zinner SH, Denny-Brown BC, Braun P, et al: Risk of infection with intravenous indwelling catheters—Effect of application of antibiotic ointment. *J Infect Dis* 120:616-619, 1969
33. Felig P: Prevention of intravenous loss of catheters. *New Eng J Med* 276:1263-1264, 1967

Primary Hyperparathyroidism

Two Cases—Carcinoma and Adenoma—In a Small Hospital

MARK J. DE MEO, M.D., *San Rafael*,
NORMAN M. SHAPIRO, M.D., AND
KAZIMIERZ W. POMYKALSKI, M.D.,
San Francisco

SINCE ASKANAZY first described an abnormal parathyroid in 1904,¹ several types of parathyroid disease have been delineated. Because of the protean symptoms associated with it, primary parathyroidism is frequently not considered in diagnosis until the clinical manifestations have completely developed or serum calcium is determined fortuitously, as it was in the two cases here presented. In both cases the patients were admitted for reasons other than primary hyperparathyroidism, and elevated serum calcium levels were discovered on routine admission Chemistry Panel 12 (Technicon SMA 12-60) determinations.

Numerous follow-up studies of primary parathyroid disease make the point that adequate correlation of surgical and pathologic findings during operation can lead to appropriate curative surgical operation.²⁻⁶ All too often this correlation is not made until after the procedure is finished, necessitating further operation later.

This presentation will emphasize routine Chemistry Panel 12 admission analyses⁷ and the close correlation of pathologic and surgical findings at the time of the first exploratory surgical procedure for primary parathyroid disease.

Reports of Cases

Case 1. A 68-year-old white woman was admitted to hospital October 1 1969 for elective repair of a right femoral hernia. Constipation and vague bone discomfort had been present for some years. On physical examination no masses were palpated in the neck. A preoperative Panel 12 (SMA 12-60) determination revealed calcium of 16 mg per 100 ml. Phosphorus was 1.9 mg per 100 ml and alkaline phosphatase was 90 millimoles per ml. Blood urea nitrogen was 17 mg per 100 ml and total protein was 6.9 grams, of which albumin was 4.5 grams. Creatinine was 1.9 mg per 100 ml

Submitted, revised, December 8, 1970.

Reprint requests to: Terra Linda Medical Laboratories, 750 Las Gallinas Avenue, San Rafael, Ca. 94903 (Dr. M. J. De Meo).




Figure 1.—Parathyroid carcinoma, showing pronounced cellular pleomorphism and trabecular pattern (x250).




Figure 2.—Parathyroid carcinoma, showing trabecular-glandular pattern with a centrally located abnormal mitotic figure (x400).

and tubular resorption of phosphate was 40 percent (normal 80 to 90 percent). Prednisone was given, 40 mg a day for 5 days. The serum calcium remained elevated at 15.6 mg per 100 ml.

Radiographs of the phalanges revealed subperiosteal resorption in the distal portions of the proximal and midphalanges.

On October 11 an exploration of the neck was performed and a large, rubbery firm, brown-tan mass was found in the left side of the neck, partially compressing the thyroid gland. The tumor was somewhat adherent to the recurrent laryngeal nerve and adjacent tissue, but no frank invasion was noted. A frozen section of the lesion was prepared and the other glands were identified and left intact.

The postoperative course was unremarkable for several days until the serum calcium dropped to 8.4 mg per 100 ml and the patient began to have signs of tetany and paranoid delusions.⁸ Treatment was begun immediately with calcium gluconate intravenously and large doses of calcium by mouth, and the serum calcium slowly reverted to normal. The patient was discharged October 19, and all paranoid delusions slowly cleared. Serum calcium remained normal for 12 months, then increased to 13.5 mg per 100 ml.

Pathologist's Report. The specimen consisted of a 5 by 4 cm irregularly shaped, rubbery firm, brown-tan mass appearing roughly encapsulated. Cut section revealed a variegated appearance with areas of yellow-tan blending with brown-tan. Fibrous trabeculae were seen streaking throughout the parenchyma. The total weight of this specimen was 13.5 grams. Microscopically (Figures 1 and 2) the tumor was consistent with para-

thyroid origin and had a decidedly variegated appearance. Some areas showed fairly regular cells with small, oval, somewhat hyperchromatic nuclei with eosinophilic cytoplasm. Most of the microscopic fields showed large, fibrous septae scattered throughout the parenchyma. These fibrous trabeculae separated the tissue into a glandular, trabecular pattern with individual cells showing pronounced pleomorphism with spindle-shaped nuclei and hyperchromatism containing numerous atypical mitotic figures.⁹ Distinct invasion of the capsule was seen in numerous sections.

Case 2. A 68-year-old white woman, a retired school teacher, was admitted to hospital October 28 1969 for evaluation of severe lower extremity cramps, increasing constipation and lethargy. No abnormality was noted on physical examination, and no masses were palpated in the neck. An x-ray film of the chest, sigmoidoscopic examination and radiographic examination of the upper and lower gastrointestinal tract were negative. Panel 12 (SMA 12-60) studies revealed serum calcium of 13 mg per 100 ml. Phosphorus content was 2.3 mg and urea nitrogen 23 mg per 100 ml. Total protein was 6.9 grams per albumin 4.2 grams, alkaline phosphatase 51 millimoles per ml. The serum creatinine was 1.1 mg per 100 ml. Tubular resorption of phosphate (TRP) was calculated at 83 percent by using the formula:

$$\text{Percent TRP} = 100 \frac{(1 - \text{VP} \times \text{SC})^{10}}{(\text{VC} \times \text{SP})}$$

where P = inorganic phosphate concentration (mg per ml)
 C = creatinine concentration (mg per ml)
 S = serum
 V = volume in ml

By the method of Chambers et al,¹⁰ calcium was infused in the amount of 15 mg per kilogram of body weight over a four-hour period. Tubular resorption was measured again and the results showed a flat response with no appreciable change in the TRP rate, indicating an autonomous secretion of parathyroid hormone. Prednisone suppression, 40 mg a day for five days, was carried out and the serum calcium remained elevated at 13.1 mg per 100 ml.

On November 7 1969, surgical exploration of the neck was carried out, and a well encapsulated, chocolate brown mass was removed from the right side of the neck. The mass appeared encapsulated and not adherent to any of the adjacent structures. A frozen section of the lesion and a biopsy specimen and frozen section of a normal parathyroid were prepared. A diagnosis of parathyroid adenoma was then submitted. The post-operative course was uneventful and the patient was discharged November 12 1969. The serum calcium levels thereafter remained within normal limits and the bone discomfort and constipation were relieved.

Pathologist's Report. The first specimen consisted of a well encapsulated brown-tan 3 by 2 cm encapsulated mass weighing 2 grams. On section this specimen was homogeneous and brown-tan. The second specimen consisted of a 0.5 cm portion of yellow-tan, rubbery firm tissue.

On microscopic examination the large mass revealed numerous small, compact cells with eosinophilic, somewhat granular cytoplasm. Numerous capillaries were seen throughout the mass. In one section adjacent to the capsule a normal rim of parathyroid tissue was seen compressed. The mass was completely encapsulated. The second specimen consisted of a regular appearing parathyroid gland with a normal amount of fat.

Discussion

The pathologic changes of primary hyperparathyroidism are of five classes: primary hyperplasia, single adenoma, combined adenoma and hyperplasia, multiple adenomas with or without the association of multiple endocrine adenomas, and carcinoma.^{2,5,9,11-16} Primary hyperplasia arising *de novo* or from an unknown stimulus is separated into the water-clear cell type described by Albright¹¹ and the chief cell hyperplasia more recently described by Cope and others.^{2,3,6} The

latter has caused much confusion for pathologists because of the similar pathologic appearance of primary chief cell hyperplasia and adenoma. The distinction between these two is, at times, difficult if not impossible when only one gland is examined. Although a functioning adenoma is the most common parathyroid lesion in patients with primary hypoparathyroidism,^{9,12,16-19} chief cell hyperplasia with or without association with multiple endocrine adenomas occurs with enough frequency to be of practical importance.^{2,9} Chief cell hyperplasia usually involves all parathyroid glands. Therefore, surgical treatment is considerably different than for parathyroid adenoma.^{3,20} Because excision of an adenoma results in cure^{21,22} and extirpation of anything less than three and one-half glands in primary chief cell hyperplasia will most likely result in continuing or recurrent hyperparathyroidism,³ it is of utmost importance that the surgeon and pathologist work closely together at the time of the first neck exploration to evaluate all parathyroid glands with biopsy of at least the one other parathyroid than that involved in the lesion.² Physical characteristics of the gland—consistency, color and shape and size—are of no appreciable aid in distinguishing between adenoma and chief cell hyperplasia. The microscopic pattern and cytological appearance are frequently very similar in adenomas and chief cell hyperplasia.¹⁴ The presence of a compressed rim of normal appearing parenchyma at the margin of an enlarged gland does give support to a diagnosis of adenoma, but is not a pathognomonic finding as it once was thought to be.² Decreased fat content and microscopic nodularity, seen so often in chief cell hyperplasia, can also be of help in the differentiation. If one gland is enlarged and adenomatous, with or without a compressed normal rim of tissue, and the remaining glands are normal or atrophic, adenoma is almost a certainty.² Differentiation by this means necessitates exposure of all four glands and biopsy of at least one other gland than the one involved in the lesion, since primary parathyroid hyperplasia usually affects the remaining glands. The possibility of multiple adenomas can be excluded through the careful identification of all glands and biopsy of any suspicious-appearing parathyroids.^{14-16,22}

Primary hyperplasia of the clear cell variety is rare. The microscopic and cytological patterns are distinct enough to cause the pathologist no great problems.^{9,12,14,23}

The problem of combined adenoma and hyperplasia has recently been studied^{5,13,15} and found to be distinguishable from the other causes of parathyroid disease by the procedure just described.

Carcinoma of the parathyroid gland is said to occur in only 3 to 4 percent of all cases of primary hyperparathyroidism.^{12,15,24} All parathyroid carcinomas are said to be associated with endocrinologic activity and well marked bone disease,^{12,15,24,25} as in Case 1 herein reported. In that case the surgeon suspected the neoplasm was malignant because of the adherence to the recurrent laryngeal nerve. Adherence to surrounding structures is often a helpful clue to malignancy in parathyroid neoplasms.^{12,26} It is usually indicative of capsular invasion, which is one of the criteria for diagnosis of parathyroid carcinoma.^{9,12} In this respect the criteria for the diagnosis of parathyroid carcinoma parallel those for carcinoma of the thyroid. The neoplasm is so rare that studies of series big enough to permit interpretation of the biological activity of this neoplasm have not been possible. It is thought that the usual spread is first through lymph channels.¹²

Microscopically, parathyroid carcinoma is different from adenoma in a number of ways. As in adenoma, the pattern is variable. Usually, however, capsular invasion is found, although that alone is not sufficient for diagnosis.^{9,12} Trabeculation and enlarged atypical mitotic figures are also significant criteria. Mitoses are absent in adenomas and present in carcinomas.^{12,24,27} Distinguishing parathyroid carcinomas from atypical adenomas may be extremely difficult. Black and Ackerman⁹ restricted the diagnosis of carcinoma to cases in which there is local invasion initially or distal metastasis. We believe that those criteria are too restrictive and that relying entirely on them may give the attending physician a feeling of overconfidence and delay proper treatment. The findings of pronounced cellular pleomorphism, mitotic activity and capsular invasion are more realistic for the diagnosis of parathyroid carcinoma.

Summary

Two cases of primary hyperparathyroidism (adenoma and carcinoma) were discovered serendipitously in a small community hospital by

Chemistry Panel 12 (Technicon, SMA 12-60) determinations. A discussion of the various types of parathyroid pathology is presented with emphasis on the correlation of gross, surgical and pathologic microscopic findings at the time of the first exploratory operation. Also stressed is the biopsy of more than one gland to differentiate chief cell hyperplasia from adenoma. The gross and microscopic criteria for the diagnosis of parathyroid carcinoma are reviewed.

REFERENCES

1. Askanazy M: *Über Ostitis deformans ohne Gewebe*. Arb a d Geb Path Anat Inst zu Tübingen 4:398-244, 1902-1904
2. Black WC, Utley J: The differential diagnosis of parathyroid adenoma and chief cell hyperplasia. *Amer J Clin Path* 49:761-775, 1968
3. Cope O, Keynes WM, Roth SI, et al: Primary chief cell hyperplasia of the parathyroid glands: A new entity in the surgery of hyperparathyroidism. *Ann Surg* 148:375-388, 1958
4. Goldsmith RS: A rapid calcium infusion test for hyperparathyroidism. *J Clin Endocr* 22:741-747, 1962
5. Golden A, Canary J, Kerwin M: Concurrence of hyperplasia and neoplasia of the parathyroid glands. *Amer J Med* 38:562-578, 1965
6. Black W, Haff RC: The surgical pathology of parathyroid chief cell hyperplasia. *Amer J Clin Path* 53:565-579, 1970
7. Boonstra CE, Jackson CE: Hyperparathyroidism detected by routine serum calcium analysis. *Ann Intern Med* 63:468, 1965
8. Vincent RG, Moore GE, Watne AL: Abnormal behavior following parathyroidectomy. *JAMA* 180:372, 1962
9. Black BK, Ackerman LV: Tumors of the parathyroid: A review of twenty-three cases. *Cancer* 3:415-444, 1950
10. Chambers EL Jr, Gordan GS, Goldman L, et al: Tests for hyperparathyroidism: Tubular reabsorption of phosphate, phosphate deprivation and calcium infusion. *J Clin Endocr* 16:1507-1521, 1956
11. Albright F, Bloomberg E, Castleman B, et al: Hyperparathyroidism due to diffuse hyperplasia of all the parathyroid glands rather than adenoma of one: Clinical studies on three such cases. *Arch Intern Med* 54:315-329, 1934
12. Castleman B: Tumors of the parathyroid glands. *Armed Forces Institute of Pathology Atlas of Tumor Pathology*, sec. 4, FAS, 15:1952
13. Harris JC, Rosenbaum D, Oliner L: Primary hyperparathyroidism with parathyroid hyperplasia and adenoma in a single gland. *Amer J Clin Path* 45:51-55, 1966
14. Roth SI: Pathology of the parathyroids in hyperparathyroidism. *Arch Path (Chicago)* 73:75-90, 1962
15. Straus FH II, Paloyan E: The pathology of hyperparathyroidism. *Surg Clin N Amer* 49:27-42, 1969
16. Woolner LB, Keating FR Jr, Black BM: Tumors and hyperplasia of the parathyroid glands: A review of the pathological findings in one hundred and forty cases of primary hyperparathyroidism. *Cancer* 5:1069-1088, 1952
17. Hellstrom J, Ivarmark BI: Primary hyperparathyroidism: Clinical and structural findings in one hundred and thirty-eight cases. *Acta Chir Scand Suppl* 294:5-89, 1962
18. Pyrah LN, Hodgkinson A, Anderson CK: Primary hyperthyroidism. *Brit J Surg* 53:245-316, 1966
19. Wermer P: Genetic aspects of adenomatosis of endocrine glands. *Amer J Med* 16:363-371, 1954
20. Block MA, Greenawald K, Horn RC Jr, et al: Involvement of multiple parathyroids in hyperparathyroidism. *Amer J Surg* 114:530, 1967
21. Goldsmith RE, Zalme E, Weinstein AS: Clinical experience with primary hyperparathyroidism: Evaluation of treatment. *Acta Endocr (Kobenhavn)* 52:221-238, 1966
22. Norris EH: The parathyroid adenoma: A study of three hundred and twenty-two cases. *Surg Gynec Obstet Suppl* 84:1-41, 1947
23. Roth SI, Manger BL: The cytology of the adenomatous, atrophic and hyperplastic parathyroid glands of man. *Arch Path Anat* 335:389-410, 1962
24. Barnes BA, Cope O: Carcinoma of parathyroid glands: Report of ten cases with endocrine function. *JAMA* 173:556, 1961
25. de Wesselow OLV, de Wardener HE: Carcinoma of the parathyroid gland with hyperparathyroidism. *Lancet* 1:820-823, 1949
26. Cope O: The study of hyperparathyroidism at the Massachusetts General Hospital. *New Eng J Med* 274:1174-1182, 1966
27. Norris EH: Carcinoma of the parathyroid glands with a preliminary report of three cases. *Int Abstr Surg* 86:1-21, 1948